

# Essential Descriptive Inorganic Chemistry

What every chemistry student should know

Peter G. Nelson



Download free books at

**bookboon**.com

PETER G. NELSON

---

# **ESSENTIAL DESCRIPTIVE INORGANIC CHEMISTRY**

WHAT EVERY CHEMISTRY  
STUDENT SHOULD KNOW

Essential Descriptive Inorganic Chemistry: What every chemistry student should know

1<sup>st</sup> edition

© 2016 Peter G. Nelson & [bookboon.com](http://bookboon.com)

ISBN 978-87-403-1502-8

Peer review by Dr David A. Johnson, emeritus reader at the Open University, UK

# CONTENTS

	<b>Introduction</b>	<b>6</b>
<b>1</b>	<b>Hydrogen</b>	<b>8</b>
<b>2</b>	<b>Oxygen</b>	<b>10</b>
<b>3</b>	<b>Carbon</b>	<b>12</b>
<b>4</b>	<b>Calcium</b>	<b>15</b>
<b>5</b>	<b>Sodium and Potassium</b>	<b>18</b>
<b>6</b>	<b>Chlorine</b>	<b>21</b>
<b>7</b>	<b>Fluorine</b>	<b>24</b>
<b>8</b>	<b>Nitrogen</b>	<b>27</b>
<b>9</b>	<b>Sulfur</b>	<b>31</b>



[www.sylvania.com](http://www.sylvania.com)

**We do not reinvent  
the wheel we reinvent  
light.**

Fascinating lighting offers an infinite spectrum of possibilities: Innovative technologies and new markets provide both opportunities and challenges. An environment in which your expertise is in high demand. Enjoy the supportive working atmosphere within our global group and benefit from international career paths. Implement sustainable ideas in close cooperation with other specialists and contribute to influencing our future. Come and join us in reinventing light every day.

Light is OSRAM

**OSRAM  
SYLVANIA** 

<b>10</b>	<b>Phosphorus</b>	<b>35</b>
<b>11</b>	<b>Silicon</b>	<b>38</b>
<b>12</b>	<b>Iron</b>	<b>40</b>
<b>13</b>	<b>Aluminium</b>	<b>44</b>
<b>14</b>	<b>Copper</b>	<b>47</b>
<b>15</b>	<b>Manganese</b>	<b>50</b>
<b>16</b>	<b>Chromium</b>	<b>52</b>
<b>17</b>	<b>Nickel</b>	<b>55</b>
<b>18</b>	<b>Zinc</b>	<b>56</b>
<b>19</b>	<b>Other Important Elements</b>	<b>57</b>
<b>20</b>	<b>Other Notable Elements</b>	<b>70</b>
	<b>Endnotes</b>	<b>73</b>

# INTRODUCTION

Descriptive inorganic chemistry is a big subject. Comprehensive texts (e.g. *Comprehensive Inorganic Chemistry*) run into several volumes. No student is capable of learning all the facts.

Educators have responded to this by arguing that what students need to do is not learn facts but the principles and theories behind the facts. The problem with this approach is that chemical principles and theories are not good enough for this. They only explain a small fraction of all the facts. Teaching principles and theories without any facts therefore leaves students ignorant of many of the facts, including very basic ones.

This short text seeks to resolve this problem. It sets out the descriptive inorganic chemistry every chemistry student should know. Only the most important chemistry of the most important elements is included. These are the elements that come high in rankings based on natural abundance, world consumption, and number of compounds.<sup>1</sup> Elements are not grouped according to the Periodic Table as this can obscure their distinctive character. Considerations of periodicity complement the material presented here.

I have, for the most part, given substances their traditional names. These are the names in, for example, the *British National Formulary*, used by doctors and pharmacists. In some cases, these differ from the names recommended by the International Union of Pure and Applied Chemistry (IUPAC). In these cases, I also give the IUPAC name. I have followed IUPAC in my spelling of 'sulfur'.

I have also, for the most part, given substances their traditional formulae. Some of these are empirical (e.g. NaCl), some molecular (e.g. HCl), some mixed (e.g. Na<sub>2</sub>O<sub>2</sub>). Ideally, these should be differentiated as in, for example, (NaCl), [HCl], and (Na<sub>2</sub>[O<sub>2</sub>]), but this is cumbersome. As a general rule, (1) volatile substances are molecular, (2) involatile substances that are soluble in nonionizing volatile solvents are 'macromolecular' (i.e. they contain large molecules like C<sub>60</sub>), and (3) involatile substances that are insoluble in nonionizing volatile solvents are 'nonmolecular' (i.e. have a framework structure like diamond or sodium chloride). Types (1) and (2) have molecular formulae, type 3 empirical formulae (or a mixed formula if they contain molecular ions).

In general formulae, M represents a metal atom, X a non-metal atom, and R a radical ( $\text{CH}_3$ ,  $\text{C}_2\text{H}_5$ , etc.). ‘Organic chemistry’ is defined as the chemistry of hydrocarbons and their derivatives.

### *Acknowledgement*


I am very grateful to Dr. David Johnson of the Open University for reading through drafts of this book and commenting extensively on them.

### *Illustrations*


I have taken the illustrations from the internet, with my thanks. I have specified the sources in the captions. As far as I know, all are free to copy, but if any are not, I apologize.

# 1 HYDROGEN


On earth, hydrogen occurs mainly as *water* ( $\text{H}_2\text{O}$ ). Rain water contains dissolved carbon dioxide from the atmosphere and is slightly acidic:



Mineral waters contain dissolved salts. Pure water is obtained by distillation. This conducts electricity slightly through self-ionization:




Hydrogen ions are hydrated [ $\text{H}(\text{H}_2\text{O})_4^+$  in dilute solution]. Acids increase the concentration of  $\text{H}^+(\text{aq})$  ions; alkalis increase the concentration of  $\text{OH}^-(\text{aq})$ . Acids turn blue litmus red, alkalis red litmus blue.






Electrolysis of water in the laboratory  
[LibreTexts]

Hydrogen ( $\text{H}_2$ ) is obtained from water by electrolysis, the water being made more conducting by the addition of an acid or alkali:






It is also prepared industrially by heating coke (from coal) or hydrocarbons (from petroleum oil) in steam:




The mixture of CO and H<sub>2</sub> is called *water gas*.

In the laboratory, hydrogen can be made by the action of a dilute acid on zinc and other metals:




The gas is colourless, odourless, and very light. It burns in air or oxygen with a pale blue flame:




Some mixtures are explosive (hydrogen in a test-tube ignites with a ‘pop’).


Hydrogen combines with other non-metals, e.g.



It also combines with some metals, e.g.



Sodium hydride is a colourless solid containing H<sup>-</sup> ions and is a useful agent for converting RCl to RH:




Large quantities of hydrogen are used to make ammonia (NH<sub>3</sub>) and to hydrogenate unsaturated fats.

Prolonged electrolysis of water leads to a separation of isotopes, deuterium oxide (D<sub>2</sub>O) accumulating in the residue.


## 2 OXYGEN

Oxygen (O<sub>2</sub>) is a colourless, odourless gas, comprising about 1/5th of the atmosphere by volume. It is isolated by fractional distillation of liquid air and kept under pressure in steel cylinders.


The gas combines with most other elements to form oxides. In these, oxygen atoms are almost always bivalent. For example, it combines with hydrogen to form water (H–O–H):



There are a great many organic compounds containing oxygen atoms. They burn in air or oxygen to form carbon dioxide and water. The reaction is more vigorous in oxygen and can be dangerous (oxygen will reignite a glowing splint). Controlled oxidation of carbohydrates takes place in the body:





The reverse process occurs in photosynthesis.




Wooden splint burning in oxygen  
[Kevin A. Boudreaux]

Sodium burns in an excess of oxygen to give a pale yellow solid which, when added to an acid, gives a solution of hydrogen *peroxide* (H–O–O–H). (The solid is mainly sodium peroxide,  $(\text{Na}^+)_2[\text{O}_2]^{2-}$ , the yellow colour coming from a trace of *superoxide*,  $(\text{Na}^+)[\text{O}_2]^-$ .)




The solution has strong oxidizing properties. It is used as a bleach and in the treatment of wastes. Pure hydrogen peroxide can be obtained by fractional distillation under reduced pressure, but the process is hazardous. The substance is made industrially by the oxidation of a quinol.

Oxygen can be prepared in the laboratory by a number of methods, e.g. by dropping water on sodium peroxide:



If an electric discharge is passed through oxygen, *ozone* is formed:



This is a blue gas with a pungent smell. It is even more reactive than oxygen.

### 3 CARBON

Carbon exists in several forms. Important ones are *graphite* and *diamond*. Both occur naturally. Impure graphite is made by heating wood or coal in the absence of air to give respectively *charcoal* or *coke*. These are used as fuels, in metallurgy, and as absorbents. Diamonds are made by heating graphite under pressure. Diamond has a 3D framework structure, graphite a 2D one. Graphite is black, and conducts electricity in 2D. Both are very involatile. Other forms include *buckminsterfullerene* ( $C_{60}$ ) and *graphene* (a single layer of graphite on a substrate).



Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)



**Deloitte.**

© Deloitte & Touche LLP and affiliated entities.





Diamond and graphite  
[Wikimedia]

Carbon burns in a limited supply of air to give carbon monoxide and in excess to give carbon dioxide:





Carbon dioxide is a colourless, odourless gas, used in carbonated drinks. It is denser than air and does not support combustion, hence its use as a fire extinguisher. On freezing, it condenses under atmospheric pressure to a white solid (*dry ice*) and under higher pressure to a colourless liquid. Both are used as refrigerants.


Carbon dioxide dissolves in water to give weakly acidic solutions:




It reacts with alkalis to give salts of  $\text{CO}_3^{2-}$  and  $\text{HCO}_3^-$  ions:



These are respectively carbonates and *bicarbonates* (hydrogencarbonates). They are colourless (unless the cation is coloured). Calcium carbonate occurs naturally (*limestone*). Carbon dioxide can be prepared in the laboratory by the action of an acid on a carbonate, e.g.




Carbon monoxide is a colourless, odourless, very poisonous gas. It reacts with some metals to give *carbonyls*, from which useful derivatives can be made:




Metal carbonyls are very toxic.

Hydrocarbons occur in natural gas and petroleum oil. Natural gas comprises mainly methane ( $\text{CH}_4$ ). The components of oil are reformed by heating (*cracking*) and separated by fractional distillation. Hydrocarbons burn in air or oxygen to carbon dioxide, e.g.



Carbon dioxide in the atmosphere affects the climate.


On heating, carbon combines with many metals to form carbides, e.g.




Iron carbide is a grey, metallic compound and is present in steel.

## 4 CALCIUM

Calcium occurs naturally in *limestone* or *chalk* (calcium carbonate,  $\text{CaCO}_3$ ). When heated, this is converted into *quicklime* or *lime* (calcium oxide,  $\text{CaO}$ ):




Quicklime reacts vigorously with water to give *slaked lime* (calcium hydroxide,  $\text{Ca(OH)}_2$ ):




All three compounds are white solids. Limestone is used to make cement by heating it with clay.

SIMPLY CLEVER

ŠKODA



We will turn your CV into an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand? We will appreciate and reward both your enthusiasm and talent. Send us your CV. You will be surprised where it can take you.

Send us your CV on [www.employerforlife.com](http://www.employerforlife.com)


Download free eBooks at [bookboon.com](http://bookboon.com)

Click on the ad to read more




Lime works  
[FreeFoto]

Calcium hydroxide is sparingly soluble in water to give an alkaline solution (*lime water*):




Bubbling carbon dioxide through the solution gives a white precipitate of calcium carbonate:



This is a test for  $\text{CO}_2$ .

Treatment of calcium hydroxide with acids gives calcium salts:




These are colourless (unless the anion is coloured), but impart a brick-red colour to a flame. Calcium hydroxide is used to reduce acidity in soils.



Calcium sulfate occurs naturally as *gypsum* ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ). This loses water on heating to give a powder that readily takes up water to form a solid mass. This is used as plaster.


Calcium chloride crystallizes as a hydrate ( $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$ ). The anhydrous salt ( $\text{CaCl}_2$ ) can be obtained by heating, and is used as a desiccant.

Calcium salts of the organic anions in soaps are insoluble, hence the scum formed when washing in hard water. Hard water produces a scale when heated because of the reaction:




The reverse reaction occurs in the cold and leads to the erosion of limestone.


Calcium itself is made by electrolysis of molten calcium chloride:



It is also be made by heating calcium oxide with aluminium (an example of the *thermit* reaction):



The element is a fairly hard, grey metal. It reacts with water to give hydrogen:



Calcium  
[Metabunk]

## 5 SODIUM AND POTASSIUM

### Sodium

Sodium occurs naturally in *rock salt* (common salt, sodium chloride, NaCl) and *brine* (aqueous NaCl).



Rock salt  
[The Spice Lab]

I joined MITAS because  
I wanted **real responsibility**

The Graduate Programme  
for Engineers and Geoscientists  
[www.discovermitas.com](http://www.discovermitas.com)



**Month 16**

I was a construction supervisor in the North Sea advising and helping foremen solve problems


Real work  
International opportunities  
Three work placements




 **MAERSK**

Important compounds are *soda-ash* (sodium carbonate,  $\text{Na}_2\text{CO}_3$ ), *sodium bicarbonate* (sodium hydrogencarbonate,  $\text{NaHCO}_3$ ), and *caustic soda* (sodium hydroxide,  $\text{NaOH}$ ). These are all white solids and used as alkalis.

Sodium is made by electrolysis of molten sodium chloride:



Sodium hydroxide is made similarly by electrolysis of aqueous sodium chloride:



Sodium bicarbonate is made by bubbling carbon dioxide through an aqueous solution of sodium chloride and ammonia (the *Solvay* process):



Ammonia is regenerated from the aqueous ammonium chloride produced by heating it with lime (made, along with the carbon dioxide used, by heating limestone):




Heating sodium bicarbonate gives the carbonate:



The carbonate is also formed by burning seaweed, hence the name ‘soda-ash’.

Sodium bicarbonate is a mild alkali because of the equilibrium:





Sodium carbonate is a stronger alkali because of two equilibria, this and:






Sodium hydroxide is a very strong alkali and can burn the skin.

All three alkalis react with acids to form sodium salts, e.g.



Sodium salts are colourless (unless the anion is coloured), but impart a characteristic yellow colour to a flame. Most are freely soluble in water. Sodium nitrate ( $\text{NaNO}_3$ ) occurs naturally as *Chile saltpetre*.

Sodium itself is a soft, grey metal. It tarnishes rapidly in air and reacts vigorously with water:



It is accordingly kept under paraffin.



Sodium metal  
[Weebly]


## Potassium

Potassium is similar to sodium. An important difference is that it is essential for plant growth. The ash from burning land plants (*potash*) is potassium carbonate ( $\text{K}_2\text{CO}_3$ ). The most important ore is *sylvite* (potassium chloride,  $\text{KCl}$ ). This is used under the name 'potash' as a fertilizer. Potassium salts impart a lilac colour to a flame.


Potassium burns in air or oxygen to the yellow superoxide ( $\text{KO}_2$ ).

## 6 CHLORINE

Chlorine (Cl<sub>2</sub>) is a greenish yellow gas with a characteristic pungent smell. It is made by electrolysis of aqueous sodium chloride (NaCl):



Sodium chloride occurs naturally as *rock salt*. The gas is stored in metal cylinders.



Chlorine  
[IndiaMART]

#1 EUROPEAN BUSINESS SCHOOL  
FINANCIAL TIMES 2013

#gobeyond

### MASTER IN MANAGEMENT


**Because achieving your dreams is your greatest challenge.** IE Business School's Master in Management taught in English, Spanish or bilingually, trains young high performance professionals at the beginning of their career through an innovative and stimulating program that will help them reach their full potential.

- Choose your area of specialization.
- Customize your master through the different options offered.
- Global Immersion Weeks in locations such as London, Silicon Valley or Shanghai.


*Because you change, we change with you.*

[www.ie.edu/master-management](http://www.ie.edu/master-management)
[mim.admissions@ie.edu](mailto:mim.admissions@ie.edu)

Hydrogen burns in chlorine to form hydrogen chloride:





This is a colourless gas with a sharp, irritating smell. It fumes in moist air and dissolves very readily in water to give hydrochloric acid:




This is a strong acid with many uses.

Chlorine combines with other elements to form chlorides. These can also be made using hydrogen chloride or hydrochloric acid. For example, iron reacts with hydrogen chloride to give its lower chloride ( $\text{FeCl}_2$ ), while chlorine gives the higher chloride ( $\text{FeCl}_3$ ):




In most chlorine compounds, the element is univalent.

Chlorine combines with hydrocarbons to form chloro-compounds:




These can also be made by treating alcohols with a hydrolysable chloride, e.g.  $\text{PCl}_5$ :



Chlorine partially dissolves in water and is used to kill bacteria in water supplies. The gas dissolves in aqueous sodium hydroxide to give a solution used as a bleach. This contains *hypochlorite* ions ( $\text{ClO}^-$ , chlorate(I)):



With calcium hydroxide, *bleaching powder* is produced:







Chlorine also forms *chlorite* ( $\text{ClO}_2^-$ , chlorate(III)), *chlorate* ( $\text{ClO}_3^-$ , chlorate(V)), and *perchlorate* ions ( $\text{ClO}_4^-$ , chlorate(VII)). Perchlorates are used when a non-complexing anion is required.

Chlorine is widely used in the manufacture of chlorinated plastics such as PVC.

Chlorine can be prepared in the laboratory by several methods, the simplest being by the action of hydrochloric acid on bleaching powder:




Hydrogen chloride can be prepared by the action of concentrated sulfuric acid on sodium chloride:




## 7 FLUORINE

The most important fluorine mineral is *fluorspar* or *fluorite* (calcium fluoride,  $\text{CaF}_2$ ). Treatment of this with sulfuric acid gives hydrogen fluoride:



This is a colourless gas which condenses to a colourless liquid at  $19^\circ\text{C}$ . It is kept in steel cylinders. Because it attacks glass and causes serious burns, it has to be handled in metal or polyfluorocarbon apparatus using protective clothing. The liquid is a very good solvent.

Hydrogen fluoride readily dissolves in water to give *hydrofluoric acid*. This can be handled in polythene. Reaction with glass gives  $[\text{SiF}_6]^{2-}$  ions:



"I studied English for 16 years but...  
...I finally learned to speak it in just six lessons"



Jane, Chinese architect

ENGLISH OUT THERE

Click to hear me talking before and after my unique course download



Hydrofluoric acid is a weak acid:




Reaction with alkalis gives salts containing colourless  $\text{F}^-$  and  $\text{HF}_2^-$  ions (fluorides and hydrogendifluorides).




Hydrofluoric acid in polythene bottle  
[Wikipedia]


Fluorine itself is obtained by electrolysis of hydrogen fluoride made conducting with potassium fluoride:



This is a pale yellow gas with a sickly smell. It is stored in steel cylinders and handled in metal or polyfluorocarbon apparatus. It reacts readily with most other elements to give fluorides. It also readily displaces other non-metals from their compounds, e.g.




For elements having more than one valency, fluorine brings out a high valency. For example, sulfur burns in fluorine to give sulfur hexafluoride ( $\text{SF}_6$ ):



Chlorine under similar conditions gives  $\text{SCl}_2$ . Fluorine brings out high valencies in some metals. For example, nickel normally has a valency of two, but heating a mixture of  $2\text{KCl}$  and  $\text{NiCl}_2$  in fluorine gives quadrivalent  $\text{K}_2\text{NiF}_6$ . In its compounds, fluorine is always univalent.


Fluorine forms many organic compounds, with some or all of the hydrogen atoms replaced by fluorine. The latter are known as *perfluoro*-compounds, and are made by fluorinating the corresponding hydrogen compounds with fluorine or a reactive fluoride. Other fluoro-compounds are made by various methods, e.g.




## 8 NITROGEN

Nitrogen ( $\text{N}_2$ ) is a colourless, odourless gas, comprising about 4/5ths of the atmosphere by volume. It is isolated by fractional distillation of liquid air and kept under pressure in steel cylinders.


Nitrogen partially combines with hydrogen when these are heated together under pressure in the presence of an iron catalyst (the *Haber* process):



The product (*ammonia*) is separated by liquefaction and kept in steel cylinders. It is a pungent gas which dissolves in water to give a weakly alkaline solution:



Addition of acids gives *ammonium* salts:



Excellent Economics and Business programmes at:



university of  
 groningen





“The perfect start  
 of a successful,  
 international career.”

**CLICK HERE**  
 to discover why both socially  
 and academically the University  
 of Groningen is one of the best  
 places for a student to be


[www.rug.nl/feb/education](http://www.rug.nl/feb/education)

These are colourless unless the anion is coloured. Ammonia can be prepared in the laboratory from ammonium chloride and slaked lime:




Liquid ammonia is a solvent. It dissolves metallic sodium and potassium to give blue solutions containing  $\text{M}^+$  ions and free electrons.


Ammonia burns in air or oxygen to nitrogen:




However, in the presence of a platinum catalyst, nitrogen monoxide (NO) is formed:



This is a colourless gas, which, in excess of air or oxygen, reacts further to give nitrogen dioxide ( $\text{NO}_2$ ):




This is a brown gas, which reacts with water to give *nitric acid*:



This can be concentrated by distilling it with concentrated sulfuric acid. The concentrated acid is a brownish liquid as a result of partial decomposition into nitrogen dioxide:





Nitric acid is a strong acid, and combines with alkalis to form *nitrates*:




These are colourless unless the cations are coloured. Sodium and potassium nitrates occur naturally ( $\text{KNO}_3$ , *nitre* or *saltpetre*;  $\text{NaNO}_3$ , *Chile saltpetre*). Ammonium nitrate ( $\text{NH}_4\text{NO}_3$ ) is used widely as a fertilizer along with ammonia itself and other derivatives.

As well as being a strong acid, nitric acid is a strong oxidizing agent, and dissolves some metals that other acids will not, e.g. copper:



Copper dissolving in concentrated nitric acid  
[Exploratorium]



Nitrogen forms a second series of salts (*nitrites*). These can be made by reducing nitrates, e.g.



Nitrogen forms other hydrides and oxides. These include *hydrazine* ( $\text{N}_2\text{H}_4$ ), made by oxidizing ammonia with aqueous sodium hypochlorite:




This is a powerful reducing agent, being readily oxidized to  $\text{N}_2$ . Among the other oxides is *nitrous oxide* or ‘laughing gas’ (dinitrogen monoxide,  $\text{N}_2\text{O}$ ), made by heating ammonium nitrate or, more safely, a mixture of ammonium sulfate and sodium nitrate:




The gas supports combustion and rekindles a glowing splint.

When ammonia is heated with methane over a platinum catalyst, *hydrogen cyanide* (HCN) is formed:



This is a colourless gas with the smell of almonds. Its solution in water is weakly acidic, and reacts with alkalis to produce *cyanides*:



These are colourless when the cations are colourless. The gas and its salts are extremely poisonous. Cyanide ions form strong complexes with some metal ions, e.g.  $[\text{Fe}(\text{CN})_6]^{4-}$ .

There are a great many organic compounds containing nitrogen. These include amines ( $\text{RNH}_2$ ,  $\text{R}_2\text{NH}$ ,  $\text{R}_3\text{N}$ ) and nitro-compounds ( $\text{RNO}_2$ ).

**American online**  
**LIGS University**  
is currently enrolling in the  
Interactive Online **BBA, MBA, MSc,**  
**DBA and PhD** programs:

- ▶ enroll **by September 30th, 2014** and
- ▶ **save up to 16%** on the tuition!
- ▶ pay in 10 installments / 2 years
- ▶ Interactive Online education
- ▶ visit [www.ligsuniversity.com](http://www.ligsuniversity.com) to find out more!

**Note:** LIGS University is not accredited by any nationally recognized accrediting agency listed by the US Secretary of Education. More info [here](#).






## 9 SULFUR

Sulfur occurs naturally in a few locations as a bright yellow crystalline solid (*brimstone*). This can be extracted by melting it (the *Frasch* process). In this form it comprises  $S_8$  molecules in which sulfur atoms are bound into an eight-membered ring. It can be obtained in other forms ( $S_n$ ). Since the atomicity varies, I shall use the empirical formula S.




Sulfur  
[Mini Me Geology]



Sulfur burns in air to sulfur dioxide:



This is a colourless gas with a choking smell. It dissolves in water to give a weakly acidic solution:




Reactions with alkalis give salts of  $SO_3^{2-}$  and  $HSO_3^-$  ions:




These are called respectively *sulfites* and *bisulfites* (hydrogensulphites). They are colourless (unless the cation is coloured) and are useful reducing agents, being readily oxidized.




Sulfur dioxide combines further with oxygen in the presence of a catalyst to give sulfur trioxide:



This is a colourless gas which condenses to a white solid containing  $(\text{SO}_3)_3$  molecules. It reacts vigorously with a stoichiometric quantity of water to give hydrogen sulfate or *concentrated sulfuric acid (oil of vitriol)*:



This is a colourless, oily liquid. It is very corrosive and burns the skin. It reacts vigorously with more water to form *sulfuric acid*:




This is a strong acid and widely used. Sulfates are colourless (unless the cation is coloured). Some occur naturally, e.g.  $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$  (*gypsum*).




Sulfuric acid plant  
 [DSD Chemtech]

Sulfur dioxide can be prepared in the laboratory by the action of concentrated sulfuric acid on a sulfite, e.g.






Sulfur forms other oxo-compounds. When a solution of a sulfite is boiled with sulfur, a *thiosulfate* is formed, analogous to  $\text{SO}_4^{2-}$ :





Sulfur burns in fluorine to give sulfur hexafluoride:



This is a colourless, odourless, unreactive gas. Its inertness makes it very useful.

On heating, sulfur combines with many metals to form sulfides, e.g.



.....Alcatel-Lucent 


[www.alcatel-lucent.com/careers](http://www.alcatel-lucent.com/careers)

What if you could build your future and create the future?

One generation's transformation is the next's status quo. In the near future, people may soon think it's strange that devices ever had to be "plugged in." To obtain that status, there needs to be "The Shift".




Iron monosulfide is a grey, nonstoichiometric solid. It reacts with acids to give hydrogen sulphide:




Iron disulfide is golden-yellow in colour and occurs naturally as *iron pyrites* or *fools' gold*. It contains  $\text{S}_2^{2-}$  ions.


Hydrogen sulfide is a colourless gas with the odour of bad eggs. It is extremely poisonous. It occurs naturally in sour natural gas, from which it can be extracted with an aqueous amine:



It can also be obtained by reducing organic sulfides in petroleum oil:




The gas is oxidized to sulfur using a catalyst (the *Claus* process):






# 10 PHOSPHORUS

Phosphorus occurs most widely as *phosphate rock*. This has the approximate composition  $\text{Ca}_3(\text{PO}_4)_2$ . Treatment of this with concentrated sulfuric acid gives hydrogen phosphate:




This is a colourless, low-melting solid, which readily dissolves in water to give *phosphoric acid*:




Concentrated solutions of this (typically 85%) are syrupy. Reaction with alkalis gives salts containing colourless  $\text{PO}_4^{3-}$ ,  $\text{HPO}_4^{2-}$ , and  $\text{H}_2\text{PO}_4^-$  ions (phosphate, hydrogenphosphate, dihydrogenphosphate). Ammonium phosphate is used as a fertilizer.

Hydrogen-containing phosphate ions can combine together by loss of water molecules to form a wide variety of polynuclear ions. For example,



The product in this case contains  $-\text{O}-\text{PO}_2^--\text{O}-\text{PO}_2^-$  chains. DNA contains similar chains.

Phosphorus itself can be obtained by heating phosphate rock with sand ( $\text{SiO}_2$ ) and coke (C):




Condensation of the vapour gives white phosphorus, a waxy solid comprising  $\text{P}_4$  molecules. Since this easily catches fire, it is kept under water. From it, red and black forms can be obtained by the application of heat and pressure. These have nonmolecular structures ( $\text{P}_n$ ). I shall represent the element by the empirical formula P.



Red and white phosphorus  
[BBC]

Phosphorus burns in an excess of air or oxygen to *phosphorus pentoxide*,  $(\text{P}_2\text{O}_5)_2$ :



In the past four years we have drilled

# 81,000 km

That's more than **twice** around the world.

**Who are we?**  
We are the world's leading oilfield services company. Working globally—often in remote and challenging locations—we invent, design, engineer, manufacture, apply, and maintain technology to help customers find and produce oil and gas safely.

**Who are we looking for?**  
We offer countless opportunities in the following domains:

- **Engineering, Research, and Operations**
- **Geoscience and Petrotechnical**
- **Commercial and Business**


If you are a self-motivated graduate looking for a dynamic career, apply to join our team.

[careers.slb.com](https://careers.slb.com)

**What will you be?**



## Schlumberger

This is a white solid that readily takes up water and is used as a drying agent:




Phosphorus also forms lower oxides and acids.


Phosphorus burns in a limited supply of chlorine to phosphorus trichloride and in excess to phosphorus pentachloride:




The trichloride is a colourless liquid, the pentachloride a white solid. With a Grignard reagent, P-Cl can be converted to P-R:



Phosphorus combines with many metals to form phosphides, e.g.



Hydrolysis of calcium phosphide gives *phosphine*:




This is a colourless gas with the smell of rotten fish and is poisonous.

# 11 SILICON

Silicon occurs naturally in rocks and sands as silicon dioxide (*silica*,  $\text{SiO}_2$ ) and metal silicates (e.g. *feldspars*). These are used as building materials. Silica is fused with alkalis (typically  $\text{Na}_2\text{CO}_3$  and  $\text{CaCO}_3$ ) to make glass. Feldspars are sodium, potassium, or calcium aluminosilicates (e.g.  $\text{NaAlSi}_3\text{O}_8$ ).

Silicon also occurs in clays. These are baked to make bricks, and with limestone ( $\text{CaCO}_3$ ) to make cement. Clays are aluminium silicates (e.g. china clay,  $\text{Al}_2(\text{OH})_4\text{Si}_2\text{O}_5$ ).

Silicon is extracted from silica by heating this with coke:




Pure silicon is an involatile lustrous grey semiconductor. It can be doped with other elements to change its electrical properties and make 'silicon chips'.




Silicon  
[WebElements]


Silicon combines with chlorine to form silicon tetrachloride:




This is a colourless liquid which fumes in moist air:




It is a starting point for the preparation of organosilicon compounds. For example, treatment with two moles of a Grignard reagent gives  $\text{R}_2\text{SiCl}_2$ :



The same product can be obtained by passing RCl over hot silicon in the presence of copper:



Hydrolysis of this gives a polymer:



This is a *silicone*. Terminal groups are made by including  $\text{R}_3\text{SiCl}$  and branching groups,  $\text{RSiCl}_3$ . Silicones have a variety of uses.

When heated with an excess of carbon, silicon forms silicon carbide ( $\text{SiC}$ ). When pure this is colourless, but the commercial material has a dark colour. It is very hard and used as an abrasive (*carborundum*).

Silicon combines with metals to form *silicides* (e.g.  $\text{Mg}_2\text{Si}$ ). Treatment of these with acid gives silicon hydrides (*silanes*,  $\text{SiH}_4$ ,  $\text{Si}_2\text{H}_6$ ,  $\text{Si}_3\text{H}_8$  etc.). These are inflammable in air.


**Maastricht University** *Leading in Learning!*

**Join the best at the Maastricht University School of Business and Economics!**

**Top master's programmes**

- 33<sup>rd</sup> place Financial Times worldwide ranking: MSc International Business
- 1<sup>st</sup> place: MSc International Business
- 1<sup>st</sup> place: MSc Financial Economics
- 2<sup>nd</sup> place: MSc Management of Learning
- 2<sup>nd</sup> place: MSc Economics
- 2<sup>nd</sup> place: MSc Econometrics and Operations Research
- 2<sup>nd</sup> place: MSc Global Supply Chain Management and Change

Sources: Keuzegids Master ranking 2013; Elsevier 'Beste Studies' ranking 2012; Financial Times Global Masters in Management ranking 2012

**Visit us and find out why we are the best!**  
**Master's Open Day: 22 February 2014**


**Maastricht University is the best specialist university in the Netherlands (Elsevier)**

**www.mastersopenday.nl**



## 12 IRON

Small quantities of iron occur naturally in meteorites. The metal is extracted from iron oxide ores (usually red-brown *haematite*,  $\text{Fe}_2\text{O}_3$ ) by heating the ore with coke (C) and limestone ( $\text{CaCO}_3$ ) in a blast of air to give *pig-iron*. The net reaction is



The limestone removes impurities from the ore as a slag. Pig-iron contains carbon in the form of iron carbide ( $\text{Fe}_3\text{C}$ ). The concentration of this is reduced by blowing oxygen through molten pig-iron in a converter to make *steel*. Other metals are added to make the kind of steel required, e.g. chromium for stainless steel.



Iron ore  
[Canadian Business Journal]






Blast furnace  
[Pinterest]


Iron is a malleable grey metal, which rusts in moist air to a red-brown powder of approximate composition  $\text{Fe}(\text{OH})_3$ :




The element forms two main series of compounds: bivalent (*ferrous* or iron(II)) and trivalent (*ferric* or iron(III)). Thus it dissolves in mineral acids to form pale green  $\text{Fe}^{2+}(\text{aq})$  ions:





These are readily oxidized to  $\text{Fe}^{3+}(\text{aq})$  ions, e.g. with hydrogen peroxide:



Aqueous  $\text{Fe}^{3+}$  ions are pale violet, but solutions containing them are often yellow because of hydrolysis or complexing:



In neutral or alkaline solutions,  $\text{Fe}^{2+}$  ions are oxidized by air. Thus addition of aqueous sodium hydroxide to a  $\text{Fe}^{2+}$  solution precipitates pale green ferrous hydroxide which darkens on exposure to air to form a red-brown, gelatinous precipitate of ferric hydroxide



Ferric hydroxide contains variable amounts of water.



**> Apply now**

REDEFINE YOUR FUTURE  
**AXA GLOBAL GRADUATE  
PROGRAM 2015**

redefining / standards 


agence c&g - © Photonostop




Ferrous sulfate,  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  (green vitriol)  
[IndiaMART]

Both  $\text{Fe}^{2+}$  and  $\text{Fe}^{3+}$  ions form complexes, e.g.  $[\text{Fe}(\text{CN})_6]^{4-}$  and  $[\text{Fe}(\text{CN})_6]^{3-}$ . The pigment *Prussian blue* is a mixed  $\text{Fe}^{\text{II}}$ - $\text{Fe}^{\text{III}}$  cyano-complex. In *haemoglobin*, an  $\text{Fe}^{2+}$  ion is coordinated by five nitrogen atoms, leaving one coordination position to bind an  $\text{O}_2$  molecule.

Iron also forms organometallic compounds. For example, powdered iron combines with carbon monoxide to give iron pentacarbonyl:



This is a toxic, yellow liquid. Treatment of anhydrous iron dichloride with sodium cyclopentadienyl in ether gives orange crystals of *ferrocene*:




This has a sandwich structure. Both compounds satisfy the 18-electron rule.




## 13 ALUMINIUM

Large quantities of aluminium occur in silicate rocks, sands, and clays. The main ore is *bauxite* ( $\text{AlO}_x(\text{OH})_{3-2x}$ ). It also occurs as *cryolite* ( $\text{Na}_3\text{AlF}_6$ ).

The metal is extracted by electrolysis of a solution of aluminium oxide ( $\text{Al}_2\text{O}_3$ ) in molten cryolite (cry.):




The oxide is made by dissolving bauxite in aqueous sodium hydroxide, removing insoluble impurities, precipitating aluminium hydroxide with acid, and heating:




The cryolite, if not mined, is made from bauxite, fluorspar ( $\text{CaF}_2$ ), and soda-ash ( $\text{Na}_2\text{CO}_3$ ).

The metal is whitish grey, light, malleable, and a good conductor of electricity. It does not rust because its oxide forms a protective layer over the surface. The metal is widely used as a structural material, e.g. in aircraft.

Aluminium burns brightly when heated in air or oxygen:




The metal abstracts oxygen from many metal oxides when it is heated with them (the *thermit* reaction), e.g.




The reaction is so exothermic as to melt the iron.


In the presence of chloride ions, aluminium dissolves in mineral acids to give aluminium salts:




The chloride ions disrupt the protective oxide film. The salts are colourless when the anions are colourless. Addition of aqueous sodium hydroxide precipitates aluminium hydroxide:



This is a white, gelatinous solid containing variable amounts of water. The solid dissolves in excess of aqueous sodium hydroxide to give aluminate ions as in the purification of bauxite:



Aluminium hydroxide is thus amphoteric. The metal itself will dissolve in hydroxide solutions:



The double salt potassium aluminium sulfate (*alum*,  $\text{KAl}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ ) is used as a mordant.



**Empowering People. Improving Business.**

BI Norwegian Business School is one of Europe's largest business schools welcoming more than 20,000 students. Our programmes provide a stimulating and multi-cultural learning environment with an international outlook ultimately providing students with professional skills to meet the increasing needs of businesses.

BI offers four different two-year, full-time Master of Science (MSc) programmes that are taught entirely in English and have been designed to provide professional skills to meet the increasing need of businesses. The MSc programmes provide a stimulating and multi-cultural learning environment to give you the best platform to launch into your career.

- MSc in Business
- MSc in Financial Economics
- MSc in Strategic Marketing Management
- MSc in Leadership and Organisational Psychology

**BI NORWEGIAN BUSINESS SCHOOL**


EFMD **EQUIS** ACCREDITED

[www.bi.edu/master](http://www.bi.edu/master)




Alum crystals  
[Natural Deodorant Store]


Aluminium burns in chlorine to give aluminium chloride:



This readily takes up water, but in its anhydrous state is used in organic chemistry to catalyse Friedel-Crafts reactions. With Grignard reagents, it forms air-sensitive organoaluminium compounds, which are also used as catalysts:





Another useful reagent is lithium aluminium hydride ( $\text{LiAlH}_4$ ) made by heating the elements together under pressure:




This is a white solid rapidly hydrolysed by moisture.

# 14 COPPER

Small quantities of copper occur naturally. Most is extracted from *iron pyrite* ( $\text{FeS}_2$ ) containing small amounts of *copper pyrite* ( $\text{CuFeS}_2$ ). This is fused with silica ( $\text{SiO}_2$ ) and a blast of air forced through it. This converts the iron to iron silicate slag and releases the copper:





The copper is purified by electrolysis:




Copper is a reddish, malleable metal. It is an excellent conductor of electricity, widely used in electrical wiring. Its surface turns brown in moist air and then green. The brown colour is due to the formation of oxide, the green to copper salts derived from acid gases ( $\text{CO}_2$ ,  $\text{SO}_2$ ) in the air. The metal forms alloys with zinc (*brass*), tin (*bronze*) and nickel (*monel metal*).



Copper only dissolves in oxidizing acids, e.g. nitric acid:



It will, however, dissolve in other acids in the presence of oxygen:





Copper forms two main series of compounds: univalent (*cuprous* or copper(I)) and bivalent (*cupric* or copper(II)). Thus when heated in air or oxygen it forms black cupric oxide ( $\text{CuO}$ ), which decomposes on stronger heating to red cuprous oxide ( $\text{Cu}_2\text{O}$ ):








$\text{Cu}_2\text{O}$  and  $\text{CuO}$  dissolve in non-complexing mineral acids to give respectively colourless  $\text{Cu}^+(\text{aq})$  and blue  $\text{Cu}^{2+}(\text{aq})$  ions:



The former rapidly disproportionate:



This does not happen in strong hydrochloric acid because the cuprous chloro-complex is stronger than the cupric:



## Need help with your dissertation?

Get in-depth feedback & advice from experts in your topic area. Find out what you can do to improve the quality of your dissertation!


Get Help Now



Go to [www.helpmyassignment.co.uk](http://www.helpmyassignment.co.uk) for more info



Indeed, the reverse reaction can take place under these conditions:




Cupric sulfate,  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  (blue vitriol)  
[nontoxicprint]

Cupric complexes include dark blue  $[\text{Cu}(\text{NH}_3)_4]^{2+}$  and the oxygen-carrier in many organisms (*haemocyanin*).

# 15 MANGANESE


Manganese is extracted from the mineral *pyrolusite* (manganese dioxide,  $\text{MnO}_2$ ) by heating this with coke (C):



It is a grey metal and used to harden steel.


The element forms compounds in various valencies. The most important of these are *manganous* (manganese(II)) salts, the dioxide, and potassium *permanganate* (manganate(VII)).

Manganous salts can be made by dissolving the metal in mineral acids:




Manganous ions are very pale pink (dilute solutions are colourless).

Manganese dioxide is a black solid, usually slightly deficient in oxygen. Potassium permanganate is made by first fusing the dioxide with potassium hydroxide in air to give potassium *manganate* (manganate(VI)):



The manganate then is oxidized to permanganate by electrolysis:




Potassium manganate is dark green, the permanganate deep purple.



Potassium permanganate crystals dissolving in water  
[Victoria State Government]

Potassium permanganate is a powerful oxidizing agent. Aqueous solutions are used to titrate reducing agents in the presence of sulphuric acid, e.g. oxalates:




Treatment of a permanganate with cold concentrated sulphuric acid gives (di)manganese heptoxide,  $\text{Mn}_2\text{O}_7$ . This is a green oil, and liable to explode violently.


Manganese forms organometallic compounds, e.g.  $\text{Mn}_2(\text{CO})_{10}$ .

# 16 CHROMIUM


Chromium is extracted from the mineral *chromite* ( $\text{FeCr}_2\text{O}_4$ ). It is a light grey metal and is used to produce stainless steel. For this purpose, the ore is heated with coke to give a chromium-iron alloy for addition to steel:



For other purposes, chromite is fused with sodium hydroxide in air and the product, sodium *chromate*, extracted from the iron oxide also produced with water:



Sodium chromate is yellow. Acid converts it into orange sodium *dichromate*:




Brain power

By 2020, wind could provide one-tenth of our planet's electricity needs. Already today, SKF's innovative know-how is crucial to running a large proportion of the world's wind turbines.

Up to 25 % of the generating costs relate to maintenance. These can be reduced dramatically thanks to our systems for on-line condition monitoring and automatic lubrication. We help make it more economical to create cleaner, cheaper energy out of thin air.

By sharing our experience, expertise, and creativity, industries can boost performance beyond expectations.


Therefore we need the best employees who can meet this challenge!

The Power of Knowledge Engineering


Plug into The Power of Knowledge Engineering.  
Visit us at [www.skf.com/knowledge](http://www.skf.com/knowledge)

**SKF**


Chromates and dichromates are used as oxidizing agents. For example, sodium or potassium dichromate in acid is used to oxidize alcohols to aldehydes:



The resulting solutions contain *chromic* (chromium(III)) ions. These are violet or, if complexed [as, e.g., in  $\text{CrSO}_4^+(\text{aq})$ ], green. Further reduction with zinc gives *chromous* (chromium(II)) ions:




These are sky-blue and readily oxidized.




Solutions of chromate and dichromate  
[ColourLex]

Treatment of a chromate or dichromate with concentrated sulfuric acid gives red crystals of chromium trioxide,  $\text{CrO}_3$ .


Reduction of anhydrous sodium dichromate with carbon gives green chromic oxide:



Heating this with aluminium gives chromium (thermit process):



Objects are chromium-plated by electrolysis of a chromic solution with a chromium anode and the object as cathode:



Chromic oxide  
[Flying Sky Industries]

Chromic ions form many complexes, e.g.  $[\text{Cr}(\text{NH}_3)_6]^{3+}$  and  $[\text{Cr}(\text{CN})_6]^{3-}$ . Chromium also forms organometallic compounds, e.g.  $\text{Cr}(\text{CO})_6$ .

What do you want to do?

No matter what you want out of your future career, an employer with a broad range of operations in a load of countries will always be the ticket. Working within the Volvo Group means more than 100,000 friends and colleagues in more than 185 countries all over the world. We offer graduates great career opportunities – check out the Career section at our web site [www.volvogroup.com](http://www.volvogroup.com). We look forward to getting to know you!



**VOLVO**  
AB Volvo (publ)  
[www.volvogroup.com](http://www.volvogroup.com)

VOLVO TRUCKS | RENAULT TRUCKS | MACK TRUCKS | VOLVO BUSES | VOLVO CONSTRUCTION EQUIPMENT | VOLVO PENTA | VOLVO AERO | VOLVO IT  
VOLVO FINANCIAL SERVICES | VOLVO 3P | VOLVO POWERTRAIN | VOLVO PARTS | VOLVO TECHNOLOGY | VOLVO LOGISTICS | BUSINESS AREA ASIA




# 17 NICKEL

Nickel occurs along with other metals mainly in sulfide ores. Separation from other metals depends on the ore. The resulting nickel sulfide ( $\sim\text{NiS}$ ) is roasted in air to the oxide and reduced with coke:




Nickel is malleable whitish-grey metal. It does not tarnish and is used in coinage. It and its alloy with copper (*monel metal*) are used to handle fluorine.

Nickel readily dissolves in dilute mineral acids to give solutions of nickel salts:



Aqueous nickel ions are green. Addition of sodium hydroxide precipitates green nickel hydroxide, which is insoluble in excess:



Oxidation gives a black precipitate of variable composition with nickel in a higher valency.





Nickel sulfate,  $\text{NiSO}_4 \cdot 7\text{H}_2\text{O}$   
[IndiaMART]

Nickel forms complexes, e.g.  $[\text{Ni}(\text{NH}_3)_6]^{2+}$  (octahedral) and  $[\text{Ni}(\text{CN})_4]^{2-}$  (square-planar). It also forms organometallic compounds, e.g. nickel carbonyl,  $\text{Ni}(\text{CO})_4$ . This is a colourless, volatile, toxic liquid. Pure nickel is obtained by passing carbon monoxide over impure metal to give the carbonyl which is then decomposed by heating:




## 18 ZINC


The main ore is *zinc blende* (zinc sulfide,  $\text{ZnS}$ ). This is roasted to the oxide and reduced with coke:




Zinc is a brittle, whitish-grey metal. It is used as a protective coating for iron (*galvanization*) and in copper-zinc alloys (*brass*). It dissolves in mineral acids to give solutions of zinc salts:




These are colourless unless the anion is coloured. Addition of aqueous sodium hydroxide precipitates white zinc hydroxide:



This is amphoteric and dissolves in excess:



Zinc itself dissolves in hot aqueous sodium hydroxide with the evolution of hydrogen;



Zinc oxide and carbonate (*calamine*,  $\text{ZnCO}_3$ ) are white and used in ointments. Zinc sulfide is also white, but fluoresces when exposed to high-energy radiation (e.g. alpha-rays). Zinc oxide is yellow when hot.





Zinc sulfate,  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$  (white vitriol)  
[Wikipedia]

# 19 OTHER IMPORTANT ELEMENTS

## Magnesium

Magnesium is similar to calcium. The chief difference is that the metal is very light, and is used for constructional purposes, e.g. in aircraft. So too are its alloys with aluminium.

A major source is *dolomite* ( $\text{MgCa}(\text{CO}_3)_2$ ). This is calcined and then heated with an iron-silicon alloy (the *Pidgeon* process):



Magnesium also occurs as magnesium chloride ( $\text{MgCl}_2$ ) in some brines and is isolated electrolytically.

Magnesium compounds include *magnesia* ( $\text{MgO}$ ), *Epsom salt* ( $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ ), and *chlorophylls* ( $\text{Mg}^{2+}$  complexes). *Milk of magnesia* is a suspension of  $\text{Mg}(\text{OH})_2$  in water.

**gaiteye**  
Challenge the way we run

EXPERIENCE THE POWER OF  
FULL ENGAGEMENT...

.....


RUN FASTER.  
RUN LONGER..  
RUN EASIER...

READ MORE & PRE-ORDER TODAY  
[WWW.GAITEYE.COM](http://WWW.GAITEYE.COM)




Epsom salt  
[Dr. Teal's]

Magnesium reacts with alkyl and aryl chlorides, bromides, and iodides in dry ether to give the Grignard reagents used widely in organic chemistry:





### **Bromine and iodine**

Bromine and iodine are similar to chlorine. Some brines contain their ions, and bromine and iodine can be extracted from them by treatment with chlorine:



Bromine ( $\text{Br}_2$ ) is a dark red-brown liquid with a red-brown vapour. Iodine ( $\text{I}_2$ ) forms shiny black crystals with a violet vapour. It dissolves in organic solvents to give violet solutions, but in aqueous potassium iodide to give a yellow-brown solution:






[A Review of the Universe]

Like hydrogen chloride, hydrogen bromide (HBr) and hydrogen iodide (HI) are colourless gases, which dissolve readily in water to form strong acids (*hydrobromic* and *hydriodic* respectively).


Fluorine, chlorine, bromine, and iodine all form salts containing  $X^-$  ions and are known collectively as *halogens*.

## Boron


Boron is similar to silicon except that it is trivalent in its compounds. It occurs as borates (e.g. *borax*,  $\text{Na}_2\text{B}_4\text{O}_5(\text{OH})_4 \cdot 8\text{H}_2\text{O}$ ). Treatment of these with hot mineral acids yields colourless crystals of boric acid on cooling. For example:




Boric acid is a weak monobasic acid:



It is used to make Pyrex glass. Heating it gives the oxide,  $\text{B}_2\text{O}_3$ :




Boron trifluoride ( $\text{BF}_3$ ) is an important catalyst. It is made by heating a borate or the oxide with fluorspar ( $\text{CaF}_2$ ) and concentrated sulfuric acid:



The trifluoride is a colourless gas and forms many adducts, e.g. with ammonia ( $\text{H}_3\text{NBF}_3$ ). Reaction with metal fluorides gives  $[\text{BF}_4]^-$  salts.

Boron trichloride ( $\text{BCl}_3$ ) is made by heating the oxide with chlorine in the presence of carbon:



This reacts with Grignard reagents to form organoboron compounds ( $\text{BR}_3$ ), used in organic synthesis.

Boron itself can be made by reducing boron compounds. The purest samples are black and, like silicon, involatile and semiconducting. They are either amorphous or crystalline, with many different structures.

This e-book  
*is made with*  
**SetaPDF**



PDF components for PHP developers

[www.setasign.com](http://www.setasign.com)






Boron  
[Lenntech]


Boron combines with metals to form *borides* (e.g.  $\text{MgB}_6$ ). Treatment of these with acid gives boron hydrides (*boranes*,  $\text{B}_2\text{H}_6$ ,  $\text{B}_4\text{H}_{10}$  etc.). Like silanes, these are inflammable in air.

## Titanium

Titanium occurs as *ilmenite* ( $\text{FeTiO}_3$ ) and *rutile* ( $\text{TiO}_2$ ). It is extracted by heating the ore with coke and chlorine, e.g.




Titanium tetrachloride is a colourless liquid and is separated from iron trichloride by distillation. It is then reduced by heating it with magnesium:



Titanium is a grey metal. It is used as a constructional material because of its high strength-to-weight ratio.

Titanium dioxide is widely used as a white pigment. One way of making it is to heat the tetrachloride in oxygen:









Titanium dioxide  
[Nanotech Etc.]

Titanium tetrachloride fumes in moist air and is used to make smoke screens:




It is also mixed with  $\text{Al}_2(\text{C}_2\text{H}_5)_6$  to catalyse the polymerization of olefins (*Ziegler-Natta* catalysis).

A solution of titanium tetrachloride in hydrochloric acid contains  $\text{Ti}(\text{OH})_2^{2+}$  ions and other species:




Reduction of this with zinc gives a violet solution containing  $\text{Ti}^{3+}$  ions:



This is used as a reducing agent.

## Cobalt

Cobalt occurs along with other metals in combination with sulfur and arsenic. The ores are roasted to oxides and separated by various methods. The resulting cobalt oxide ( $\text{Co}_3\text{O}_4$ ) is reduced to the metal by heating it with coke:



The metal is bluish grey and used in steel and magnets. It remains ferromagnetic to a higher temperature than other elements.

The chemistry of cobalt is similar to that of iron. Like the latter, it forms two main series of compounds: bivalent (*cobaltous* or cobalt(II)) and trivalent (*cobaltic* or cobalt(III)). Aqueous  $\text{Co}^{2+}$  ions are pink,  $\text{Co}^{3+}$  blue. The latter are unstable, being readily reduced by water:



Cobaltous sulfate,  $\text{CoSO}_4 \cdot 7\text{H}_2\text{O}$   
[Prayug Agro]

[www.sylvania.com](http://www.sylvania.com)

We do not reinvent  
the wheel we reinvent  
light.

Fascinating lighting offers an infinite spectrum of possibilities: Innovative technologies and new markets provide both opportunities and challenges. An environment in which your expertise is in high demand. Enjoy the supportive working atmosphere within our global group and benefit from international career paths. Implement sustainable ideas in close cooperation with other specialists and contribute to influencing our future. Come and join us in reinventing light every day.


Light is OSRAM

Both ions form complexes.  $\text{Co}^{3+}$  complexes are more stable than the aqueous ion and are particularly numerous, interconversions taking place only very slowly. For example, treatment of  $\text{CoCl}_2(\text{aq})$  with ammonia in the presence of air gives five different compounds depending on the conditions: golden-brown  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$ , bright red  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{O})]\text{Cl}_3$ , purple  $[\text{CoCl}(\text{NH}_3)_5]\text{Cl}_2$ , deep violet *cis*- $[\text{CoCl}_2(\text{NH}_3)_4]\text{Cl}$  and green *trans*- $[\text{CoCl}_2(\text{NH}_3)_4]\text{Cl}$ . These were used by Werner to develop his coordination theory. Most  $\text{Co}^{3+}$  complexes are octahedral. Vitamin  $\text{B}_{12}$  is a  $\text{Co}^{3+}$  complex.


Cobalt forms organometallic compounds. The simplest carbonyl is  $\text{Co}_2(\text{CO})_8$ .

## Silver


Small amounts of silver occur naturally. Most is obtained from the slime that is formed under the anode in the electrolytic purification of copper. Silver is isolated from this chemically and purified by electrolysis in dilute nitric acid:



Silver is a malleable, whitish grey metal and good conductor of electricity (the best of all the elements). It dissolves in dilute nitric acid to give a solution of silver nitrate:



This is colourless. Silver chloride (white), bromide (pale yellow), and iodide (yellow) are insoluble and darken on exposure to light:




Adding silver nitrate to sodium chloride  
[BBC]

Silver forms complexes, e.g.  $[\text{Ag}(\text{NH}_3)_2]^+$  and  $[\text{Ag}(\text{CN})_2]^-$ .



## Gold and platinum

These occur naturally. Gold is a yellow metal, platinum whitish grey.

The two metals do not dissolve in ordinary acids, but they do dissolve in *aqua regia* (a mixture of concentrated nitric and hydrochloric acids). The resulting solutions contain respectively  $[\text{AuCl}_4]^-$  and  $[\text{PtCl}_6]^{2-}$  ions:



Gold also dissolves in aerated sodium cyanide solution, a process used to extract it from crushed rock:



Gold forms two main series of compounds, *aurous* (gold(I)) and *auric* (gold(III)). Platinum likewise forms *platinous* (platinum(II)) and *platinic* (platinum(IV)) compounds. Auric and platinous complexes ( $d^8$ ) are characteristically square-planar.




Chloroauric acid,  $\text{H}[\text{AuCl}_4] \cdot 3\text{H}_2\text{O}$   
[Alibaba]



Platinum is an important catalyst. It is used, for example, in catalytic converters in cars. Complexes of gold and platinum are used in medicine.

## Tin

The most important ore is *cassiterite* ( $\text{SnO}_2$ ). This is reduced by heating it with coke:



The metal is whitish grey and has a relatively low melting point ( $232^\circ\text{C}$ ). It forms two series of compounds: bivalent (*stannous*, tin(II)) and quadrivalent (*stannic*, tin(IV)), e.g.



Stannous chloride is a white solid, stannic chloride a colourless liquid.



Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)

**Deloitte.**

© Deloitte & Touche LLP and affiliated entities.






Stannous chloride,  $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$   
[SafestChina]

Stannous compounds are reducing agents. Aqueous  $\text{Sn}^{2+}$  ions are oxidized by air.


Tin forms important alloys with copper (*bronze*) and lead (*solder*), and is used to provide a protective cover for steel (*tin-plate*) used in canning. Molten tin is used in the manufacture of float glass.

## Lead

The most important ore is *galena* ( $\text{PbS}$ ). This is roasted in a limited supply of air:



The oxide is then reduced with coke:



Lead a dense, grey, malleable metal. It is similar to tin in forming two series of compounds, bivalent (*plumbous*, lead(II)) and quadrivalent (*plumbic*, lead(IV)), e.g. colourless  $\text{Pb}^{2+}$  salts and brown  $\text{PbO}_2$ . The relative stabilities are, however, different: plumbic compounds are oxidizing agents; aqueous  $\text{Pb}^{2+}$  ions are stable in air.




Plumbous nitrate,  $\text{Pb}(\text{NO}_3)_2$   
[IndiaMART]


Lead-tin alloys are low-melting and used as solder.

### Uranium


This occurs in oxide minerals (e.g. *pitchblende*,  $\text{U}_3\text{O}_8$ ) dispersed in rocks. The rocks are crushed, leached with sulfuric acid, and the uranium isolated chemically. The resulting 'yellow cake' has the approximate composition  $\text{U}_3\text{O}_8$  and, on one scheme, is reduced by heating it with hydrogen:



The dioxide is converted to the tetrafluoride by heating it in hydrogen fluoride:




The tetrafluoride is reduced by heating it with magnesium:



[Uranium Mining Museum]




Natural uranium comprises mainly  $^{235}\text{U}$  (0.7%) and  $^{238}\text{U}$  (99.3%), the first of which is used in nuclear reactors. To separate these, the tetrafluoride is converted to the hexafluoride by heating it in fluorine:




The hexafluoride is volatile and is separated in a gas centrifuge,  $^{238}\text{UF}_6$  concentrating towards the wall and  $^{235}\text{UF}_6$  towards the centre. Chemists use 'depleted uranium' made by removing the  $^{235}\text{UF}_6$  and reducing the  $^{238}\text{UF}_6$ .

Other sexivalent compounds include the oxide,  $\text{UO}_3$ , and salts of the yellow *uranyl* ion,  $\text{UO}_2^{2+}$ .

SIMPLY CLEVER



We will turn your CV into an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?  
We will appreciate and reward both your enthusiasm and talent.  
Send us your CV. You will be surprised where it can take you.

Send us your CV on  
[www.employerforlife.com](http://www.employerforlife.com)

## 20 OTHER NOTABLE ELEMENTS

### Lithium

This is similar to sodium and potassium. An important compound is lithium aluminium hydride,  $\text{LiAlH}_4$ , used as a reducing and hydrogenating agent in organic chemistry. Lithium salts are used in psychiatric medicine.

Lithium, sodium, and potassium, along with rubidium and caesium, are known collectively as *alkali metals*. They all react with water giving hydrogen and form basic hydroxides ( $\text{MOH}$ ) and colourless  $\text{M}^+(\text{aq})$  ions. They are distinguished by their flame colours: lithium, crimson-red; sodium, yellow; potassium, lilac; rubidium, ruby-red; caesium, sky-blue.

### Barium

This is similar to calcium. An important compound is barium sulfate ( $\text{BaSO}_4$ ). This is insoluble in water, and the formation of a white precipitate on addition of  $\text{Ba}^{2+}(\text{aq})$  is used as a test for sulfate. Being dense, barium sulfate is given to patients when X-raying the stomach and intestine.

Calcium, strontium, barium, and radium are known collectively as *alkaline earth metals*. They all react with water giving hydrogen and form basic hydroxides  $[\text{M}(\text{OH})_2]$  and colourless  $\text{M}^{2+}(\text{aq})$  ions. Their flame colours are: calcium, brick-red; strontium, crimson-red; barium, apple-green; radium, red-purple. Radium is radioactive.

### Selenium

This is similar to sulfur. Like sulfur, it forms compounds in which it has a valency of two ( $\text{H}_2\text{Se}$ ), four ( $\text{SeO}_2$ ), and six ( $\text{H}_2\text{SeO}_4$ ). It exists in several forms, from red ( $\text{Se}_8$ ) to black ( $\text{Se}_n$ ).



Black and red selenium  
[Wikipedia]

## Molybdenum and tungsten

These are whitish grey metals with high melting points. Their chemistry is dominated by oxo-compounds (*molybdates* and *tungstates*) in which they have a valency of six. Molybdenum trioxide is white (yellow when hot), tungsten trioxide yellow. Tungsten carbide (WC) is used to tip cutting tools. Both elements form organometallic compounds, e.g.  $\text{Mo}(\text{CO})_6$  and  $\text{W}(\text{CO})_6$ .

## Mercury

This is the only metal that is a liquid at room temperature. It dissolves other metals to form *amalgams*. It forms two series of compounds: *mercurous* (mercury(I), e.g.  $\text{Hg}_2\text{Cl}_2$ ) and *mercuric* (mercury(II), e.g.  $\text{HgCl}_2$ ). Aqueous  $\text{Hg}_2^{2+}$  and  $\text{Hg}^{2+}$  ions are colourless. Mercurous chloride (*calomel*) is insoluble in water. Mercury compounds are poisonous.



Mercury  
[Mining Chemicals Trading Support]

## Helium, neon, argon, krypton, and xenon

These are minor components of the atmosphere. All are monatomic. Chemically, they are almost completely or totally inactive. They are known as the *inert gases* (or, ineptly, 'noble gases'). Argon is the most abundant, comprising about 1% of the atmosphere by volume.

## Questions

- Write a short account of the chemistry of (a) calcium, (b) nitrogen, (c) iron.
- Compare and contrast the chemistry of (a) magnesium and zinc, (b) phosphorus and sulfur.
- X is a non-metal. It combines with hydrogen to form a hydride. This dissolves in water to give an acid. The acid forms a series of salts. These give a pale yellow precipitate with silver nitrate. Identify X.
- M is a metal. It forms two series of salts with valencies two and three.  $[M(H_2O)_6]^{3+}$  ions oxidize water,  $[M(NH_3)_6]^{3+}$  ions do not. Identify M.
- Which of the following represent the highest oxides of iron, manganese, and chromium?
  - $FeO_3$ ,  $MnO_2$ ,  $CrO_3$
  - $FeO_3$ ,  $Mn_2O_7$ ,  $CrO_3$
  - $Fe_2O_3$ ,  $MnO_2$ ,  $CrO_3$
  - $Fe_2O_3$ ,  $Mn_2O_7$ ,  $CrO_3$
  - $Fe_2O_3$ ,  $MnO_2$ ,  $Cr_2O_3$
  - $Fe_2O_3$ ,  $Mn_2O_7$ ,  $Cr_2O_3$

## Answers

- See the sections on these elements in the text.
- (a) Both are bivalent metals, but zinc hydroxide is amphoteric, whereas magnesium hydroxide is basic.  
(b) Their chemistries are similar, but their valencies are different (phosphorus: 3, 5; sulfur: 2, 4, 6).
- Bromine
- Cobalt
- (d). Iron does, however, form derivatives of  $FeO_3$ , e.g.  $K_2FeO_4$ , made by suspending  $Fe(OH)_3$  in  $KOH(aq)$  and oxidizing it with chlorine.

# ENDNOTES

1. See my discussion in *Introduction to Inorganic Chemistry*, [www.bookboon.com/en/introduction-to-inorganic-chemistry-ebook](http://www.bookboon.com/en/introduction-to-inorganic-chemistry-ebook), and *J. Chem. Educ.*, 1991, **68**, 732–737.

I joined MITAS because  
I wanted **real responsibility**

The Graduate Programme  
for Engineers and Geoscientists  
[www.discovermitas.com](http://www.discovermitas.com)



**Month 16**

I was a construction supervisor in the North Sea advising and helping foremen solve problems

Real work  
International opportunities  
Three work placements



 **MAERSK**